



E-ISSN: 2709-9385

P-ISSN: 2709-9377

JCRFS 2022; 3(2): 37-40

© 2022 JCRFS

www.foodresearchjournal.com

Received: 19-05-2022

Accepted: 22-06-2022

Upasana KolDepartment of Entomology,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India**Vikas Singh**Department of Entomology,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India**Gajendra Chandrakar**Department of Entomology,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India

Screening of germplasm of mungbean against pod borer complex

Upasana Kol, Vikas Singh and Gajendra Chandrakar

Abstract

The present investigation was conducted at the Research cum Instructional farm of the Indira Gandhi Krishi Vishwavidyalaya in Raipur (C.G.), during *Kharif* 2021–2022. In studies on screening of germplasm of mungbean against pod borer complex, the minimum pod damage infestation by *H. armigera* and *M. vitrata* were observed in germplasm OBG 109 and BM - 4, respectively. The highest grain yield of mungbean was recorded in KM 2241.

Keywords: Screening, mungbean, pod borer, spotted pod borer

Introduction

Mungbean is the third most important pulse crop farmed in India among the major pulse crop accounting for roughly 16% of the country's total pulse area. The pulse crop *Vigna radiata* (L.) Wilczek (Family: Leguminosae, Sub-family: Papilionaceae), also known as green gram (Synonyms: Golden bean or mung bean or moong bean), has been grown since antiquity. It is a tiny green circular shaped bean widely cultivated throughout Asia, including India, Pakistan, Bangladesh, Sri Lanka, Thailand, Laos, Cambodia, Vietnam, Indonesia, Malaysia, and the Republic of Formosa. Mungbean is a native of India and Central Asia, according to Vavilov (1926) [1]. It is a great source of high-quality protein that's easy to digest. It has a protein content of roughly 25%, which is almost three times that of grains. It meets the protein needs of the country's vegetarian population. Green gram has a nutritional content of 334 calories per 100 grams, crude protein (percentage) 24, fat (percentage) 1.3, carbohydrate (percentage) 56.6, calcium (mg / 100 g) 140, phosphorus (mg / 100 g) 280, iron (mg / 100 g) 8.4, vitamin (mg / 100 g), B1 0.47, B2 0.39, and niacin 2.0 (Baldev *et al.*, 2000) [13]. It is planted on around 4.5 million hectares, yielding 2.5 million tonnes with a productivity of 548 kg/ha and accounting for 10% of total pulse production. Green gram output is expected to reach 2.64 million tonnes in 2020 - 21, according to the Government of India's third advance projections. Green gram consumption was 22.5 lakh tonnes in the marketing year 2020 - 21, compared to production of 21.42 lakh tonnes, with the remaining demand-supply gap filled by importing roughly 1.08 lakh tonnes and opening inventories of 2.10 lakh tonnes (Green gram outlook report, AMIC, 2021). Mungbean is preyed upon by a variety of insect pests. Insect pests that attack the mungbean plant can be categorised based on how they arrive in the field, which is linked to the phenology of the plant. Stem feeders, foliage feeders, pod feeders, and storage pests are the four types. This classification is useful for determining the economic importance of pests, particularly their impact on seed yield, as well as developing control strategies. Different insect pests attack mungbean, but sucking insect pests (aphids, jassid, leaf hoppers, and whiteflies) are the most common (Islam *et al.*, 2008) [7]. Though many options are available for the management of these insect pests, farmers in India mostly use synthetic chemicals because of their quick effect with or without knowing the ill effects of these chemicals. However, farmer education for the safe and timely use of the insecticides is very important. Host plant resistance is recognised as an important component of integrated pest control since it is a low-cost, realistic, long-term strategy for sustaining lower whitefly populations and decreasing crop losses (Bellotti and Arias, 2001) [3]. Host plant resistance, according to Dowell (1990) [4], is the best long-term solution for controlling whiteflies.

Correspondence

Upasana KolDepartment of Entomology,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India

Materials and Methods

The field experiment was conducted during *Kharif* season of 2021 - 2022, at the Research cum Instructional Farm, Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G.). Total mungbean germplasm used were seventy-six, sown in RBD design with 2 replications. The observations were recorded as (i) pod damage (%): Pods damaged due to different pod borers based on the nature of damage of *Helicoverpa armigera* cause large round and regular holes on the pods, while *Maruca vitrata* makes irregular scrapping and holes on the mungbean pods was separated from 100 randomly collected pods at the time of harvest. (ii) Yield parameters: Grain yield will be recorded at the time of harvest. Afterward, the total number of pods and the number of damaged pods by pod borers on each demarcated plant were counted and converted into percentages. The percentage of pod damaged and grain yield Kg/ha were estimated with the help of following formula:

$$\text{Pod Damage (\%)} = \frac{\text{Number of damage pods}}{\text{Total Number of pods (Healthy + Damage)}} \times 100$$

$$\text{Grain yield(kg/ha)} = \frac{\text{Weight of grains in kg/plot}}{\text{Plot area in m}^2} \times 10,000$$

Statistical analysis

The data obtained were analysed statistically after using appropriate transformation. The larval population of pod borer complex data obtained was converted into square root transformation; by using the formula ($\sqrt{x} + 0.5$) the data on pod and grain damage was first recorded from the plants and then converted into percentage. The percentage data was processed under arcsine transformation $\text{Sin}^{-1}(\sqrt{x}/100)$ before statistical analysis. This transformed data was then analysed by the method of analysis of variance as described by Gomez and Gomez (1984) [5]. The "F" test was used at 5 percent level of significance. The following formulae were used for standard error, critical difference and coefficient of variance estimations:

$$C.D. = \sqrt{\frac{2EMSR}{R}} \times t (D.F. \text{ at } 5\%)$$

Results and Discussion

Screening of germplasm of mungbean against gram pod borer, *Helicoverpa armigera* (Hubner): The incidence of insect pest was assessed in the percentage pod damage at the harvesting stage of the crop. Germplasm differed significantly in terms of per cent pod damage, which ranged from 0.5 to 10.5 per cent. Among the tested germplasm, the minimum pod damage by *H. armigera* was observed in germplasm OBGG 109 with 0.5 per cent, which was found

at par with LGG 460 and PM 1711 with 1.00 per cent and IPM 1603-3, IPM 20-1, MH 1830, ML 2506, ML 818, SML 2015 with 1.5 per cent pod damage respectively, whereas the maximum pod damage was observed in germplasm BCM 20-9 with 10.5 per cent pod damage. Our findings are similar with the findings of Yadav *et al.* (2021) [12], who estimated the minimal percentages of pod infestations by *H. armigera* in the four genotypes KU-99-05, Azad Urd-1, Shekhar - 2, and PU-6 to be 5.83, 6.17, 8.50, and 9.83 per cent, respectively. Similar to this, Banu *et al.* (2007) [2] tested fifteen germplasm lines for resistance to *H. armigera* and discovered that ICPL -13201 had the lowest pod damage (25%) and greatest yield (60.35 g / plant), followed by ICPL - 13028 (28%) and ICPL-11964(29%).

Screening of germplasm of mungbean against spotted pod borer, *Maruca vitrata* (Geyer):

The insect pest incidence was observed in terms of per cent of pod damage at the harvesting stage of the crop. The germplasm showed significant differences with each other for per cent pod damage, which varied from 1.00% to 14.00%. Among the tested germplasm, the minimum pod damage by *M. vitrata* was observed in germplasm BM-4 with 1.00 per cent, which was found at par with ML 2500 at 1.00 per cent pod damage, MH 1142 with 2.5 per cent pod damage, BCM 18-1, IPM 1604 - 1, IPMD 1603 - 7, Kopergoan at 2.5 per cent pod damage, respectively. Whereas the maximum pod damage was observed in OBGG 104 with 14.00 percent. More or less the present findings are also agreement with the findings of Singh and Singh (2014) [9] who tested the 30 mungbean genotypes against *Maruca vitrata*, in which RVSm - 11 - 92 had the least pod damage, followed by PM-306 - 6, IEM 2K - 15 4, MH - 805, BM - 4, DGGS - 4, and BM - 2002 - 2. Similarly, Sandhya Rani *et al.* (2014) [8] found the preference of *M. vitrata* in five genotypes, where KM - 9 - 128, KM - 9 - 136, RMG - 492, LGG - 527, and LGG-538 were found to be tolerant, while the susceptibility of the other twenty-one genotypes ranged from 12.59 per cent (MGG - 332) to 20.0 percent (IPM - 02 - 03 and LGG - 522) and thirteen genotypes were highly susceptible ranging from 43.25 per cent (KM - 8 - 662). The others were extremely vulnerable, with a range of 20.21 per cent (UPM - 99 - 3) to 40.0 percent (KM - 2241). Srivastava and Singh (2017) [10] also found that, KM 2348 had the highest proportion of pod damage by *M. vitrata*, followed by BM 2012 - 9 (16.51%), AKM 12 - 17 (16.40%), PM 4 (15.06%) and IPM 312-20 (15.06%). Based on per cent pod damage and the Pest Susceptibility Rating (PSR) score, VGG 10 - 008 was shown to be the genotype that was least susceptible to the spotted pod borer.

Table 1: Per cent pod damage by pod borer complex and grain yield in different germplasm of mungbean (Kharif 2021-22)

S.no	Germplasm	Pod Damage (%)		Grain Yield (Kg/ha)
		<i>H. armigera</i>	<i>M. vitrata</i>	
1	AKM 12-28	6 (14.12)	6 (14.17)	806.67
2	AKM 8802	4.5 (12.07)	4.5 (12.22)	946.67
3	AKM 4	8.5 (16.88)	3 (9.83)	1110.00
4	BCM 18-1	8 (16.39)	2.5 (9.04) ^c	1016.67
5	BCM 18-2	4.5 (12.22)	10.5 (18.85)	1133.34
6	BCM 20-9	10.5 (18.85)	4 (10.53)	996.67
7	BM 2019-10	7 (15.29)	9.5 (17.93)	1036.67
8	BM-4	4 (11.44)	1 (4.06)	910.00
9	COGG 18-17	2.5 (9.04)	7.5 (15.81)	1000.00
10	COGG 912	3 (9.83)	5.5 (12.78)	1120.00
11	Daftri Vikas	6.5 (14.67)	8 (16.3)	1140.00
12	GJM 1701	12 (20.24)	11 (19.34)	1006.67
13	IPM 13-6	4 (11.44)	2.5 (9.04) ^c	970.00
14	IPM 1603-3	1.5 (6.93) ^b	12 (20.19)	1093.33
15	IPM 1604-1	3.5 (10.75)	13 (21.07)	730.00
16	IPM 20-1	1.5 (6.93) ^b	13.5 (21.51)	1043.34
17	IPM 20-2	4.5 (12.22)	8.5 (16.93)	890.00
18	IPM 2-14	3.5 (10.52)	3.5 (10.75)	973.34
19	IPM 2-3	2.5 (9.04)	5.5 (13.43)	1230.00
20	IPM 2K-14-9	2.5 (9.04)	10.5 (18.85)	1043.33
21	IPM 410-3	2.5 (8.63)	6 (14.12)	970.00
22	IPM 512-1	3.5 (10.75)	7.5 (15.81)	1133.34
23	IPMD 1603-7	6 (13.97)	2.5 (9.04) ^c	1170.00
24	JLPM 702-1	5 (12.85)	5.5 (13.43)	1016.67
25	K 851	3.5 (10.75)	7 (15.18)	1126.67
26	KM 2241	3 (9.83)	6 (14.12)	1343.34
27	KM 2419	2 (7.85)	4.5 (12.22)	1180.00
28	KM 2421	3.5 (10.52)	12.5 (20.60)	1193.33
29	Kopergoan	5(12.85)	2.5 (9.04) ^c	1036.67
30	LGG 450	3 (9.83)	9 (17.42)	1016.67
31	LGG 460	1 (5.73) ^a	4 (11.44)	1133.33
32	LGG 610	2.5 (8.63)	4 (11.53)	1000.00
33	MH 1830	1.5 (6.93) ^b	10 (18.40)	916.67
34	MH 1142	3.5 (10.52)	2.5 (8.63) ^b	1260.00
35	MH 1468	2.5 (9.04)	6 (14.12)	956.67
36	MH 1772	3.5 (10.75)	5.5 (13.43)	1246.67
37	MH 1857	6 (14.12)	4.5 (12.22)	886.67
38	MH 2-15	4 (11.44)	3.5(10.52)	866.67
39	MHBC 20-2	2.5 (9.04)	4 (11.44)	1236.67
40	MI 181-1	3 (9.83)	8 (16.3)	1110.00
41	MI 750-1	2 (8.12)	7.5 (15.87)	1170.00
42	MI 98-64	3 (9.83)	7 (15.29)	1140.00
43	ML 2500	2.5 (9.04)	1 (5.73) ^a	1066.67
44	ML 2506	1.5 (6.93) ^b	5 (12.91)	1193.34
45	ML 818	1.5 (6.93) ^b	4.5 (12.07)	1070.00
46	NVL 1143	3 (9.83)	7 (15.18)	1010.00
47	OBGG 104	2.5 (9.04)	14 (21.91)	1006.67
48	OBGG 105	3 (9.83)	9 (17.02)	1096.67
49	OBGG 109	0.5 (2.86)	3.5 (10.75)	980.00
50	OUM 11-5	3.5 (10.75)	9.5 (17.79)	1070.00
51	Palamapur 93	3 (9.83)	10.5 (18.77)	1106.67
52	PM 1711	1 (5.73) ^a	13.5 (21.51)	1020.00
53	PM 1723	5 (12.91)	11 (19.28)	1093.34
54	PM 4	4 (11.44)	7.5 (15.81)	1143.34
55	PM 6	2 (8.12)	7.5 (15.87)	1073.34
56	Pusa 0672	5(12.91)	5.5 (13.19)	1080.00
57	Pusa 1371	4.5 (12.22)	7.5 (15.81)	833.34
58	Pusa BM 9	5.5 (13.43)	4 (11.44)	950.00
59	Pusa M 2071	4.5 (12.22)	4 (11.53)	913.33
60	Pusa M 2171	6 (13.97)	6.5 (14.49)	1033.34
61	Pusa M 2172	5 (12.85)	4.5 (12.07)	1220.00
62	RM 03-71	6 (13.97)	7 (15.29)	1066.67
63	RMG 1132	4 (11.44)	6 (14.12)	1086.67
64	RMG 1139	2 (8.12)	6 (14.17)	1146.67

65	RMG 1166	4 (11.44)	5 (11.59)	1166.67
66	RVSM 18-1	5 (12.85)	3.5 (10.52)	1076.67
67	SKNM 1705	5.5 (13.54)	5.5 (13.54)	1210.00
68	SKNM 1904	4 (11.44)	5.5 (13.43)	1030.00
69	SML 1839	3 (9.97)	7 (14.98)	1246.67
70	SML 2015	1.5 (6.93) ^b	6.5 (14.67)	1036.67
71	T-44	3 (9.83)	9 (17.34)	1106.67
72	VBN 4	4.5 (12.07)	4.5 (12.07)	1046.67
73	VGG 17-049	2 (7.85)	5.5 (13.19)	1070.00
74	VGG 17-106	5.5 (13.54)	6 (13.97)	1123.34
75	VGG 18-021	3 (9.83)	3 (9.97)	1130.00
76	Pusa Vishal	3.5 (10.75)	7 (15.18)	1110.00
CD at 5% SE(m)		4.214 1.493	5.579 1.976	208.211 73.757

Figure in parenthesis () are angular transformed value

Grain Yield: The grain yield of various mungbean genotypes ranged from 730.00 kg / ha to 1343.34 kg / ha. The genotype KM 2241 was produced the highest grain yield as 1343.34 kg / ha, followed by genotype MH 1142 as 1260.00 kg / ha. Whereas genotype IPM 1604 - 1 was produced the lowest grain yield as 730.00 kg / ha, followed by genotype IPM 20 - 2 as 890.00 kg / ha. More or less the present findings are also agreement with the findings of Srivastava and Singh (2017) ^[10] who discovered that the lowest grain yield was recorded from KM 2348 (416 kg / ha), as compared to check cultivar, HUM - 12 (590 kg / ha), and that the maximum grain yield was recorded from VGG 10-008 (819 kg / ha), which was substantially different from other genotypes. Singh and Singh (2014) ^[9] also observed that AKM-4 had the highest yield, which was then followed by KM - 2293, AKM - 09 - 2, IPM - 3066, and ML - 1628.

Conclusion

On numerous metrics such as percent pod damage and grain yield, the germplasm screening trail revealed considerable differences among examined germplasm. The least pod damage impact on the germplasm OBBG 109 and BM - 4 was caused by *Helicoverpa armigera* and *Maruca vitrata*, respectively. The germplasm KM 2241 produced the highest grain yield, whereas the germplasm IPM 1604 - 1 produced the lowest grain yield.

References

1. Kebede ZY. Case study report on improved mung bean technology demonstration under North-West Ethiopia. Int. J Agric. Food Sci. 2022;4(1):35-37. DOI: 10.33545/2664844X.2022.v4.i1a.63
2. Banu RM, Muthiah AR, Ashok S. Field screening and evaluation of pigeon pea genotypes against pod borers (*Helicoverpa armigera*). Pakistan Journal of Biological Sciences. 2007;10(7):1149-1150.
3. Bellotti AC, Arias B. Host plant resistance to whiteflies with emphasis on cassava as a case study, Crop Protection. 2001;20(9):813-823.
4. Dowell R. Integrating biological control of whiteflies into crop management systems, In: D. Gerling (ed.): Whiteflies: Their Bionomics, Pest Status and Management, Intercept, Andover, U.K; c1990. p. 315-335.
5. Gomez KA, Gomez AA. Statistical procedure for Agricultural Research. John wiley and sons' publication. 2nd Edition; c1984.
6. Green gram Outlook report submitted by Agricultural Market Intelligence Centre, ANGRAU, Lam; c2021.
7. Islam MS, Latif MA, Ali M, Hossain MS. Population

- dynamics of whitefly on some recommended mungbean varieties in Bangladesh and its impact on incidence of mungbean yellow mosaic virus disease and yield, International Journal of Sustainable Agriculture Technology. 2008;5(4):41-46.
8. Rani CS, Rao RG, Chalam MSV, Kumar AP, Rao SV. Field Screening of greengram genotypes against *Maruca vitrata* in summer. Journal of Agriculture and Crop Science. 2014;1(2):18-25.
9. Singh SK, Singh PS. screening of mungbean (*Vigna radiata*) genotypes against major insects. Current Advances in Agricultural Sciences. 2014;6(1):85-87.
10. Srivastava C, Singh S. Field screening of some green gram [*Vigna radiata* (L.) Wilczek] genotypes against spotted pod borer, *Maruca vitrata* (Fabricius). Journal of Entomology and Zoology Studies. 2017;5(4):1161-1165.
11. Vavilov NI. Studies on the origin of cultivated plants. Bulletin of Applied Botany and Plant Breeding. 1926; 16(2):1-248.
12. Yadav A, Singh G, Yadav A, Singh H, Singh V, Singh p. Screening of black gram genotypes against major pod borers. Legume research - an international journal; c2021. 10.18805/LR-4686
13. Baldev B, Ramanujam S, Jain HK. Pulse Crops. Oxford IBH Publishing Company Pvt. Ltd., New Delhi; c2000. p. 229-258.